

Please visit <https://www.evidentlybetter.org/bulletins/suicide-prevention/> to view our webpage featuring key links and emerging reports about suicide prevention.

Studies

[Suicide and self-harm in nurses and midwives: urgent attention needed according to new systematic review](#)

Summarises a recent systematic review exploring the prevalence, risk factors and interventions for suicide and self-harm in nurses and midwives.

[Accessing psychological therapies following self-harm: qualitative survey of patient experiences and views on improving practice](#)

Findings identify long waiting times and inadequate service provision as barriers to high-quality and safe aftercare for patients who have self-harmed. Consistent with clinical guidelines, all patients should receive prompt aftercare and access to tailored psychological treatments following a self-harm episode.

Statistics

[Quarterly suicide death registrations in England](#)

Provisional rate and number of suicide deaths registered in England per quarter. Includes 2001 to 2021 registrations and provisional data for Quarter 1 (Jan to Mar) to Quarter 4 (Oct to Dec) 2022.

Conferences

[SSHARE NoW 6th Annual Conference](#)

The sixth annual Suicide and Self Harm Research North West conference brings together clinicians, individuals with lived experience, researchers, and academics in the field. The event is on 16 June from 9.30am to 4.30pm.

[National Confidential Inquiry into Suicide and Safety in Mental Health 9th Conference](#)

The National Confidential Inquiry into Suicide and Safety in Mental Health (NCISH) 9th NCISH Conference - Wednesday 26 April 2023 (10.00am to 12.30pm GMT). This is a free virtual conference.

[Suicide Bereavement UK's 12th International Conference](#)

One day conference: 28th September 2023 – 9am to 4.30pm (registration from 8am) Mercure Manchester Piccadilly Hotel, Manchester FACE2FACE £165 + Booking Fee + VAT LIVESTREAM £99 + Booking Fee + VAT. This event is organised to inform and guide professionals who come into contact with those bereaved by suicide and those in a position to influence policy and practice. Therefore, it is presented in a factual manner and is not suitable for people bereaved by suicide during the early stages of loss and/or struggling to cope with their loss.